


- (21) Application No 8224155
(22) Date of filing 23 Aug 1982
(30) Priority data
(31) 8125791
(32) 24 Aug 1981
(33) United Kingdom (GB)
(43) Application published
9 Mar 1983
(51) INT CL³
C07D 487/04
A61K 31/53 //
C07D 233/92
(C07D 487/04 235/00
257/00)
(52) Domestic classification
C2C 1410 142X 200 214
215 220 221 225 226 22Y
247 250 252 25Y 280 305
30Y 313 314 31Y 321 32Y
332 339 342 34Y 351 352
364 36Y 574 584 620 621
624 625 62X 631 652 662
670 671 761 802 80Y AA
KJ KR KS KT ML
U1S 1313 1314 C2C
(56) Documents cited
None
(58) Field of search
C2C
(71) Applicants
May and Baker Limited,
(Great Britain),
Dagenham,
Essex
(72) Inventors
Edward Lunt
Malcolm Francis Graham
Stevens
Robert Stone
Kenneth Robert Harry
Wooldridge
(74) Agents
J. A. Kemp and Co.,
14 South Square,
Gray's Inn,
London WC1R 5EU

(54) Tetrazine derivatives

(57) [3H] Imidazo[5,1-d] - 1,2,3,5-tetrazine - 4 - one derivatives of the formula:-




[wherein R¹ represents hydrogen, or an alkyl, alkenyl or alkynyl group containing up to 6 carbon atoms, each such group being unsubstituted or substituted by from one to three substituents selected from halogen atoms, alkoxy, alkylthio, alkylsulphinyl and alkylsulphonyl groups containing up to 4 carbon atoms, and optionally substituted phenyl groups, or R¹ represents a cycloalkyl group containing from 3 to 8 carbon atoms, and R² represents a carbamoyl group optionally N-substituted by one or two groups selected from alkyl and alkenyl groups containing up to 4 carbon atoms, and cycloalkyl groups containing 3 to 8 carbon atoms], are new therapeutically useful compounds possessing antineoplastic and immunomodulatory activity.

SPECIFICATION

Tetrazine derivatives

5 This invention relates to new [3H] - imidazo - [5,1-d] - 1,2,3,5 - tetrazin - 4 - one derivatives, to processes for their preparation and to pharmaceutical compositions containing them.

10 The compounds of the present invention are the [3H] - imidazo[5,1-d] - 1,2,3,5 - tetrazin - 4 - one derivatives of the general formula:-



15 wherein R¹ represents a hydrogen atom, or a straight- or branched-chain alkyl, alkenyl or alkynyl group containing up to 6 carbon atoms, each such group being unsubstituted or substituted by from one to three substituents selected from halogen (i.e. bromine, iodine or, preferably, chlorine or fluorine) atoms, straight- or branched-chain alkoxy, (e.g. methoxy), alkylthio, alkylsulphinyl and alkylsulphonyl groups containing up to 4 carbon atoms, and optionally substituted phenyl groups, or R¹ represents a cycloalkyl group, and R² represents a carbonyl group which may carry on the nitrogen atom one or two groups selected from straight- and branched-chain alkyl and alkenyl groups, each containing up to 4 carbon atoms, and cycloalkyl groups, e.g. a methylcarbamoyl or dimethylcarbamoyl group.

20 When the symbol R¹ represents an alkyl, alkenyl or alkynyl group substituted by two or three halogen atoms, the aforesaid halogen atoms may be the same or different. When the symbol R¹ represents an alkyl, alkenyl or alkynyl group substituted by one, two or three optionally substituted phenyl groups the optional substituents on the phenyl radical(s) may be selected from, for example, alkoxy and alkyl groups containing up to 4 carbon atoms (e.g. methoxy and/or methyl group(s)) and the nitro group; the symbol R¹ may represent, for example, a benzyl or *p*-methoxybenzyl group. Cycloalkyl groups within the definitions of symbols R¹ and R² contain 3 to 8, preferably 6, carbon atoms.

25 Preferred tetrazine derivatives of general formula I are those wherein R¹ represents a straight- or branched-chain alkyl group containing from 1 to 6 carbon atoms optionally substituted by one or two halogen (preferably chlorine, fluorine or bromine) atoms or by an alkoxy group containing 1 to 4 carbon atoms (preferably methoxy) or by a phenyl group (optionally substituted by one or two alkoxy groups containing from 1 to 4 carbon atoms, preferably methoxy), or R¹ represents an alkenyl group containing 2 to 6 carbon atoms (preferably allyl) or a cyclohexyl group.


30 More particularly preferred tetrazine derivatives are those of general formula I wherein R¹ represents a straight- or branched-chain alkyl group containing from 1 to 6 carbon atoms, and more especially from 1 to 3 carbon atoms, unsubstituted or substituted by

More especially R¹ represents a methyl or 2-haloalkyl, e.g. 2-fluoroethyl or, preferably, 2-chloroethyl, group.


35 Preferably R² represents a carbamoyl group or a monoalkylcarbamoyl, e.g. methylcarbamoyl, or monoalkenylcarbamoyl group.

40 The present invention also includes salts of the compounds of general formula I wherein R¹ represents a hydrogen atom and R² is as hereinbefore defined, more especially alkali metal, e.g. sodium, salts, and whenever the context so permits reference to the compounds of general formula I in this specification is meant to include reference to the said salts. The salts are particularly useful as intermediates.

45 According to a feature of the present invention, the compounds of general formula I, wherein R² is as hereinbefore defined and R¹ is other than hydrogen, are prepared by the reaction of a compound of the general formula:-




50 (wherein R² is as hereinbefore defined) with an isocyanate of the general formula:-



55 wherein R³ represents an alkyl, alkenyl or alkynyl group, optionally substituted by one to three substituents selected from halogen atoms, alkoxy, alkylthio, alkylsulphinyl and alkylsulphonyl groups and optionally substituted phenyl groups, or represents a cycloalkyl group, within the definition of R¹ hereinbefore recited. The reaction may be effected in the absence or presence of an anhydrous organic solvent, for example a chlorinated alkane, e.g. dichloromethane, or ethyl acetate, acetonitrile, *N*-methylpyrrolid-2-one or, preferably, hexamethylphosphoramide, at a temperature between 0° and 70°C, e.g. at the ambient temperature. The reaction may be continued for up to 30 days. Light should preferably be excluded from the reaction mixture.

60 According to a further feature of the present invention, the compounds of general formula I, wherein R² is as hereinbefore defined and R¹ is other than hydrogen, are prepared by the reaction of a compound (within general formula I) of the general formula:-



65 (wherein R² is as hereinbefore defined) or an alkali metal, e.g. sodium, salt thereof with a compound of the general formula:-



70 wherein R³ is as hereinbefore defined, and X represents the acid residue of a reactive ester, for example a halogen (e.g. chlorine) atom, or a sulphuric or sulphonic ester residue, e.g. a methoxysulphonyloxy, methanesulphonyloxy, or toluene-*p*-sulphonyloxy


represents a haloalkyl, haloalkenyl or haloalkynyl group, the acid residue of a reactive ester represented by X will be selected from those known to be not less reactive than the halogen atom substituent in R³. When X in a compound of general formula V represents a halogen atom, an alkali metal salt of the compound of general formula IV is preferably used and wherein X in a compound of general formula V represents a halogen atom and R³ is a haloalkyl, haloalkenyl or haloalkynyl group wherein the halogen atom is the same as that represented by X, an excess of the dihalo compound of general formula V is preferably used. The reaction of a compound of general formula IV or alkali metal salt thereof with a compound of general formula V, wherein R³ and X are as hereinbefore defined, may be carried out in a suitable anhydrous inert organic solvent, for example dichloromethane, acetonitrile or *N*-methylpyrrolid-2-one or mixtures thereof, at a temperature of from 0°C to 120°C and, when a compound of general formula IV is used, in the presence of an acid-binding agent, for example an alkali metal, e.g. sodium or potassium, carbonate or bicarbonate.

As a further feature of the invention, compounds of general formula IV (i.e. compounds of general formula I wherein R¹ represents a hydrogen atom and R² is as hereinbefore defined) or alkali metal salts thereof are prepared by the reaction of a compound of general formula II with a compound of the general formula:-



VI

wherein R⁴ represents an alkali metal (e.g. sodium) atom or a protecting group such as a benzyl or *p*-methoxybenzyl group, followed, when R⁴ represents a protecting group, by the replacement of the protecting group by a hydrogen atom in the compound thus obtained of the general formula:-



VII

wherein R² is as hereinbefore defined, and R⁵ represents a protecting group such as a benzyl or *p*-methoxybenzyl group, by methods known *per se*. Reaction of a compound of general formula II with a compound of general formula VI wherein R⁴ represents a protecting group may be effected as hereinbefore described for the reaction of a compound of general formula II with a compound of general formula III. Reaction of a compound of general formula II with a compound of general formula VI, wherein R⁴ represents an alkali metal atom, may be effected in a suitable inert organic solvent, e.g. ethanol, acetonitrile or *N*-methylpyrrolidone, optionally in the presence of an acid, at a temperature of from 0° to 120°C. The group R⁵ of compounds of general formula VII, wherein R² is as hereinbefore defined, may be replaced by a hydrogen atom by methods known *per se* to give a compound of general formula IV.

Compounds of general formula II may be prepared by the application or adaptation of methods known *per se*, for example methods described by Shealy

J. Org. Chem. (1961), 26, 2396.

Compounds of general formulae III, V and VI may be prepared by the application or adaptation of methods known *per se*.

By the term 'methods known *per se*' as used in the present specification is meant methods heretofore used or described in the literature.

The new tetrazine derivatives of general formula I possess valuable antineoplastic activity, for example against carcinomas, melanomas, sarcomas, lymphomas and leukaemias. They have proved particularly active in mice at daily doses between 0.5 and 16 mg/kg animal body weight, administered intraperitoneally, against TLX(S) lymphoma according to the procedure of Gescher *et al*, Biochem. Pharmacol. (1981), 30, 89, and ADJ/PC6A and M5076 (reticulum cell sarcoma). Against leukaemia L1210, grafted intraperitoneally, intracerebrally and intravenously, and P388, according to the procedure described in "Methods of Development of New Anti-cancer Drugs" (NCI Monograph 45, March 1977, pages 147-149, National Cancer Institute, Bethesda, United States), the compounds were active both intraperitoneally and orally at doses of between 2.5 and 10 mg/kg animal body weight. Inhibition of both primary tumour and metastasis was obtained against the Lewis lung carcinoma by similar dosage regimes. Against the B16 melanoma and C38 tumour in mice (NCI Monograph 45, *op cit.*) the compounds were active intraperitoneally at doses of between 6.25 and 25 mg/kg animal body weight.

The tetrazine derivatives also possess valuable immunomodulatory activity and are of use in the treatment of organ grafts and skin grafts and in the treatment of immunological diseases.

Important individual compounds of general formula I include the following:-

- 8-carbamoyl-3-methyl-[3H]-imidazo [5,1-d]-1,2,3,5-tetrazin-4-one A,
 - 8-carbamoyl-3-n-propyl-[3H]-imidazo [5,1-d]-1,2,3,5-tetrazin-4-one B,
 - 8-carbamoyl-3-(2-chloroethyl)-[3H]-imidazo [5,1-d]-1,2,3,5-tetrazin-4-one C,
 - 3-(2-chloroethyl)-8-methylcarbamoyl-[3H]-imidazo [5,1-d]-1,2,3,5-tetrazin-4-one D,
 - 8-carbamoyl-3-(3-chloropropyl)-[3H]-imidazo [5,1-d]-1,2,3,5-tetrazin-4-one E,
 - 8-carbamoyl-3-(2,3-dichloropropyl)-[3H]-imidazo [5,1-d]-1,2,3,5-tetrazin-4-one F,
 - 3-allyl-8-carbamoyl-[3H]-imidazo [5,1-d]-1,2,3,5-tetrazin-4-one G,
 - 3-(2-chloroethyl)-8-dimethylcarbamoyl-[3H]-imidazo [5,1-d]-1,2,3,5-tetrazin-4-one H,
 - 3-(2-bromoethyl)-8-carbamoyl-[3H]-imidazo [5,1-d]-1,2,3,5-tetrazin-4-one I,
 - 3-benzyl-8-carbamoyl-[3H]-imidazo [5,1-d]-1,2,3,5-tetrazin-4-one J,
 - 8-carbamoyl-3-(2-methoxyethyl)-[3H]-imidazo [5,1-d]-1,2,3,5-tetrazin-4-one K,
 - 8-carbamoyl-3-cyclohexyl-[3H]-imidazo [5,1-d]-1,2,3,5-tetrazin-4-one L,
 - and 8-carbamoyl-3-(*p*-methoxybenzyl)-[3H]-imidazo [5,1-d]-1,2,3,5-tetrazin-4-one. M
- Compounds A and D, and especially C, are of par-

The letters A to M are allocated to the compounds for easy reference later in the specification.

The following Examples illustrate the preparation of compounds of general formula I according to the present invention, and the Reference Example thereafter illustrates the preparation of intermediates.

EXAMPLE 1

Compound A

4[5]-Diazoimidazole-5[4]-carboxamide (500 mg) was suspended in methyl isocyanate (3.0 ml) and stirred in the dark, at ambient temperature, for 21 days. The reaction mixture was then diluted with anhydrous diethyl ether and filtered. The residue was washed quickly with anhydrous methanol, then with anhydrous diethyl ether, and dried in air, in the dark, at ambient temperature, to give 8-carbamoyl-3-methyl-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one, in the form of a light brown microcrystalline solid (198 mg), m.p. 210°C (with effervescence and darkening from 160° to 210°C). [Elemental analysis:—found C,36.8; H,3.10; 44.2%; C₆H₆N₆O₂ requires: C,37.1; H,3.09; N,43.3%].

EXAMPLE 2

Compound B

4[5]-Diazoimidazole-5[4]-carboxamide (300 mg) was suspended in anhydrous dichloromethane (10 ml) and treated with an excess of n-propyl isocyanate. The reaction mixture was then stirred in the dark, at ambient temperature, for 30 days. The reaction mixture was then filtered and the residue was washed quickly with anhydrous diethyl ether, and dried in air, in the dark, at ambient temperature, to give 8-carbamoyl-3-n-propyl-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one (102 mg), in the form of a pale pink powder, m.p. 167°C (with effervescence). [Elemental analysis:—found: C,43.4; H,4.57; N,38.0%; C₈H₁₀N₆O₂ requires: C,43.2; H,4.53; N,37.8%].

EXAMPLE 3

Compound C

4[5]-Diazoimidazole-5[4]-carboxamide (300 mg) was suspended in anhydrous dichloromethane (10 ml) and 2-chloroethyl isocyanate (1.0 ml) was added. The reaction mixture was then stirred in the dark, at ambient temperature, for 30 days. The cream-coloured suspension thus obtained was filtered and the residue was washed quickly with anhydrous diethyl ether and dried in air, in the dark, to give 8-carbamoyl-3-(2-chloroethyl)-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one (483 mg), in the form of a cream-coloured powder, m.p. 158°C (with vigorous decomposition). [Elemental analysis:—found: C,34.7; H,3.01; N,34.9%; C₇H₇ClN₆O₂ requires: C,34.7; H,2.91; N,34.7%].

Repetition of the above procedure has also given 8-carbamoyl-3-(2-chloroethyl)-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one in another polymorphic form, m.p. 164-165°C (with decomposition).

EXAMPLE 4

Compound A

A suspension of 4[5]-diazoimidazole-5[4]-carboxamide (1.37 g) in ethyl acetate (20 ml) was treated with methyl isocyanate (7.0 g) and was stirred

in a closed vessel in the dark at room temperature for 3 weeks. The resulting solid was filtered off and washed with diethyl ether to give 8-carbamoyl-3-methyl-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one (1.9 g), in the form of a cream-coloured solid, m.p. 212°C (with effervescence).

This material was recrystallised from three different solvent systems to give three different products, each of which had a slightly different IR spectrum.

The three products were probably all polymorphs of 8-carbamoyl-3-methyl-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one.

(i) Colourless needles were obtained from a 3:1 v/v mixture of acetone and water, ν_{\max} 3410, 3205, 1758, 1730 and 1678 cm⁻¹, m.p. 212°C (with effervescence).

(ii) White microcrystals were obtained from a 1:3 v/v mixture of acetone and water, ν_{\max} 3430, 3200, 1740 and 1675 cm⁻¹, m.p. 210°C (with effervescence).

(iii) A granular solid was obtained from hot water, ν_{\max} 3450, 3380, 3200, 1742, 1688 and 1640 cm⁻¹, m.p. 215°C (with effervescence) (darkening from 200°C).

EXAMPLE 5

Compound B

A suspension of 4[5]-diazoimidazole-5[4]-carboxamide (1.37 g) in acetonitrile (20 ml) was treated with n-propyl isocyanate (6.5 g) and was stirred in a closed vessel in the dark at room temperature for 3 weeks. The resulting pink solid was filtered off, washed with diethyl ether, and recrystallised from a mixture of water and acetone (1:4 v/v), to give 8-carbamoyl-3-n-propyl-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one (1.6 g), m.p. 170-172°C (with effervescence). By concentration of the recrystallisation mother liquor there was obtained a further quantity (0.2 g) of the same product.

EXAMPLE 6

Compound C

A suspension of 4[5]-diazoimidazole-5[4]-carboxamide (1.0 g) in ethyl acetate (30 ml) was treated with 2-chloroethyl isocyanate (3.3 ml) and the mixture was stirred in the dark, at ambient temperature, for 6 days. The reaction mixture was then diluted with diethyl ether and the resulting solid was filtered off, to give 8-carbamoyl-3-(2-chloroethyl)-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one (1.6 g) in the form of a colourless solid, m.p. 164-165°C (with decomposition). [Elemental analysis:—found: C,34.5; H,2.88; N,34.5; Cl,14.6%; C₇H₇ClN₆O₂ requires: C,34.65; H,2.91; N,34.65; Cl,14.61%].

EXAMPLE 7

Compound C

A suspension of 4[5]-diazoimidazole-5[4]-carboxamide (5.0 g) in a mixture of dichloromethane (158 ml) and N-methylpyrrolid-2-one (8.3 ml) was treated with 2-chloroethyl isocyanate (16.7 ml) and the mixture was stirred in the dark at ambient temperature for 14 days. The reaction mixture was then diluted with anhydrous diethyl ether and the resulting solid was filtered off and washed with diethyl ether, to give 8-carbamoyl-3-(2-chloroethyl)-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one (6.3 g), in the form of a purple-tinged solid, m.p. 164-165°C (with decomposition). [Elemental

analysis:— found: C,34.7; H,2.95; N,34.5; Cl,14.4%;
C₇H₇ClN₆O₂ requires: C,34.65; H,2.91; N,34.65;
Cl,14.61%].

EXAMPLE 8

5 Compound C

A suspension of 4[5] - diazoimidazole - 5[4] - carboxamide (145 g) in ethyl acetate (2175 ml) was treated with 2-chloroethyl isocyanate (478.5 ml) and stirred at 30°C, with the exclusion of light, for 2 days.

- 10 The mixture was then filtered to give 8 - carbamoyl - 3 - (2 - chloroethyl) - [3H] - imidazo[5,1-d] - 1,2,3,5 - tetrazin - 4 - one (250 g), in the form of a peach-coloured solid, m.p. 166°C.

EXAMPLE 9

15 Compound A

A stirred suspension of 4[5] - diazoimidazole - 5[4] - carboxamide (2.2 g) in a mixture of dichloromethane (70 ml) and *N* - methylpyrrolid - 2 - one (3.5 ml) was treated with methyl isocyanate (7.0 ml) and stirred at ambient temperature for 4 weeks. The mixture was diluted with diethyl ether and the resulting solid was filtered off, to give 8 - carbamoyl - 3 - methyl - [3H] - imidazo[5,1-d] - 1,2,3,5 - tetrazin - 4 - one (2.38 g), in the form of a pale purple solid, m.p. 202-203°C (with decomposition). [Elemental analysis:— found: C,36.8; H,2.94; N,43.1%; C₆H₆N₆O₂ requires: C,37.11; H,3.14; N,43.3%].

- 20 A polymorphic form of 8 - carbamoyl - 3 - methyl - [3H] - imidazo[5,1-d] - 1,2,3,5 - tetrazin - 4 - one was obtained by dissolving it in acetonitrile, filtering, concentration of the filtrate to dryness, and trituration of the resulting residue with diethyl ether. This material was in the form of an orange-tinged solid, m.p. about 200°C (with decomposition). [Elemental analysis: C,37.4; H,3.26; N,43.5%]. Its NMR spectrum in dimethylsulphoxide - D₆ was identical to that of the abovementioned pale purple solid, but its IR spectrum (KBr disc) showed some differences.

EXAMPLE 10

40 Compound D

A stirred solution of sodium nitrite (0.64 g) in water (4.6 ml) was cooled to 5°-10°C and treated dropwise at that temperature with a solution of 5 - amino - 4 - methylcarbamoylimidazole (1.00 g) in aqueous acetic acid (1M; 14.3 ml) during 5 minutes. Stirring was continued at 5°-10°C for 5 minutes. The dark red solution was then extracted with ethyl acetate (4 x 35 ml) and the combined extracts were dried over magnesium sulphate. The resulting solution contained crude 4[5] - diazo - 5[4] - methylcarbamoylimidazole, which was unstable and was used immediately for the next stage without further purification.

- The solution of 4[5] - diazo - 5[4] - methylcarbamoylimidazole in ethyl acetate, prepared as described above, was treated with 2-chloroethyl isocyanate (4.3 ml) and was allowed to stand in the dark for 1 day. The solution was then evaporated at 40°C/10 mm Hg and the residue was triturated with petroleum ether (b.p. 40°-60°C) to give an orange gum (4.23 g). This gum was treated with ethyl acetate (50 ml) and filtered, and the filtrate was evaporated at 40°C/10 mm Hg to give an orange gum (2.94 g). This gum was purified by medium pressure column chromatography on silica gel, eluting with a

give 3 - (2 - chloroethyl) - 8 - methylcarbamoyl - [3H] - imidazo[5,1-d] - 1,2,3,5 - tetrazin - 4 - one (0.81 g), in the form of a purple solid, m.p. 120-122°C (with decomposition). [Elemental analysis:— found: C,37.3; H,3.58; N,31.9%; C₈H₇ClN₆O₂ requires: C,37.4; H,3.53; N,32.7%].

EXAMPLE 11

Compound E

A suspension of 4[5] - diazoimidazole - 5[4] - carboxamide (1.0 g) in ethyl acetate (50 ml; dried over anhydrous potassium carbonate) was treated with 3-chloropropyl isocyanate (4.86 g) and the mixture was stirred at ambient temperature for 3 days. The reaction mixture was then diluted with anhydrous diethyl ether and the resulting solid was filtered off, and washed with anhydrous diethyl ether, to give 8 - carbamoyl - 3 - (3 - chloropropyl) - [3H] - imidazo[5,1-d] - 1,2,3,5 - tetrazin - 4 - one (1.05 g) in the form of a pink solid, m.p. 153-154°C (with decomposition). [Elemental analysis:— found: C,37.1; H,3.42; N,32.7; Cl,13.8%; C₈H₉ClN₆O₂ requires: C,37.4; H,3.53; N,32.8; Cl,13.8%].

EXAMPLE 12

Compound F

- 90 By proceeding in a manner similar to that described hereinbefore in Example 11 but replacing the 3-chloropropyl isocyanate used as a starting material by the appropriate quantity of 2,3 - dichloropropyl isocyanate, there was prepared 8 - carbamoyl - 3 - (2,3 - dichloropropyl) - [3H] - imidazo[5,1-d] - 1,2,3,5 - tetrazin - 4 - one, in the form of an off-white solid, m.p. 153-155°C (with decomposition). [Elemental analysis:— found: C,32.7; H,2.51; N,28.7; Cl,24.1%; C₈H₈Cl₂N₆O₂ requires C,33.0; H,2.77; N,28.9; Cl,24.4%].

EXAMPLE 13

Compound G

- Stirred allyl isocyanate (4.5 ml, redistilled immediately before use) was treated with 4[5] - diazoimidazole - 5[4] - carboxamide (1.0 g) and then with hexamethylphosphoramide (20 ml). The mixture was stirred at ambient temperature in the dark for 18 hours and then it was diluted with anhydrous diethyl ether and filtered. The resulting colourless solid was washed with anhydrous diethyl ether, to give 3 - allyl - 8 - carbamoyl - [3H] - imidazo[5,1-d] - 1,2,3,5 - tetrazin - 4 - one (1.6 g), in the form of a colourless solid, m.p. 149-150°C [ν_{\max} (KBr disc): 1730, 1675 cm⁻¹; NMR in DMSO-d₆: singlets at 8.75, 7.67 and 7.60 δ ; double double triplet at 6.02 δ (J=5.5, 8, 10Hz), double doublet at 5.35 δ (J=1.5, 8 Hz) and 5.20 δ (J=1.5, 10 Hz) and doublet at 4.88 δ (J=5.5)].

EXAMPLE 14

Compound H

- 120 A solution of 4[5] - diazo - 5[4] - dimethylcarbamoylimidazole (1.59 g; prepared as described in Reference Example 1 hereafter) in dry ethyl acetate (57 ml) was treated with 2-chloroethyl isocyanate (6.36 g) and stirred at room temperature in the dark for 24 hours. The solution was then evaporated *in vacuo* at 35°C, finally at 0.1 mm Hg to remove the excess of 2-chloroethyl isocyanate. The residual liquid was purified by medium pressure column chromatography of silica gel, eluting with a mixture

(2-chloroethyl)-8-dimethylcarbamoyl-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one (0.82 g), in the form of colourless crystals, m.p. 114-116°C. [Elemental analysis:—found: C,39.7; H,3.95; N,30.8%; C₉H₁₁ClN₆O₂ requires: C,39.9; H,4.10; N,31.0%].

EXAMPLE 15

Compound I

A stirred suspension of 4[5]-diazoimidazole-5[4]-carboxamide (1.0 g) in hexamethylphosphoramide (4 ml) was treated with 2-bromoethyl isocyanate (4.5 ml) and the mixture was stirred in the dark, at ambient temperature, for 2 days. The reaction mixture was then diluted with anhydrous diethyl ether and the resulting solid was filtered off, and washed with anhydrous diethyl ether, to give 3-(2-bromoethyl)-8-carbamoyl-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one (1.17 g), in the form of a colourless solid, m.p. 156-157°C (with decomposition). [Elemental analysis:—found: C,29.5; N,2.36; Br,27.3%; C₇H₇BrN₆O₂ requires: C,29.3; H,2.46; N,29.3; Br,27.8%].

EXAMPLE 16

Compound J

By proceeding in a manner similar to that described hereinbefore in Example 15 but replacing the 2-bromoethyl isocyanate used as a starting material by the appropriate quantity of benzyl isocyanate, there was prepared 3-benzyl-8-carbamoyl-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one (0.83 g), in the form of a buff-coloured solid, m.p. 176-177°C (with decomposition). [Elemental analysis:—found: C,53.6; H,3.66; N,31.0%; C₁₂H₁₀N₆O₂ requires: C,53.3; H,3.73; N,31.1%].

EXAMPLE 17

Compound K

A suspension of 4[5]-diazoimidazole-5[4]-carboxamide (0.3 g) in acetonitrile (5 ml) was treated with 2-methoxyethyl isocyanate (0.5 g) and the mixture was stirred at between 45° and 47°C in the dark for 24 hours. The resulting solid was filtered off and washed with diethyl ether, to give crude 8-carbamoyl-3-(2-methoxyethyl)-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one (0.45 g), m.p. 145-147°C (with decomposition).

The product was purified by recrystallisation from aqueous acetone to give pink rosettes, or from aqueous dimethylsulphoxide to give colourless needles, m.p. 164-165°C (with decomposition). [Elemental analysis: C,40.4; H,4.20; N,35.2%; C₈H₁₀N₆O₃ requires: C,40.34; H,4.20; N,35.2%].

EXAMPLE 18

Compound L

A suspension of 4[5]-diazoimidazole-5[4]-carboxamide (0.30 g) in acetonitrile (10 ml) was treated with cyclohexyl isocyanate (1.0 g) and the mixture was stirred at 60°C in the dark for 3 days. The resulting solid was filtered off and washed with a mixture of ethanol and 0.880 aqueous ammonia (100:0.5 v/v; 20 ml) for one minute, to give 8-carbamoyl-3-cyclohexyl-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one (0.015 g), m.p. 196°C (with effervescence).

EXAMPLE 19

Compound J

boxamide (0.4 g) in acetonitrile (10 ml) was treated with benzyl isocyanate (0.6 g) and the mixture was stirred at 60°C in the dark overnight. The reaction mixture was then cooled and filtered, to give 3-benzyl-8-carbamoyl-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one (0.75 g), in the form of a pale pink solid, m.p. 187-188°C (with effervescence).

EXAMPLE 20

Compound M

A suspension of 4[5]-diazoimidazole-5[4]-carboxamide (0.1 g) and *p*-methoxybenzyl isocyanate (0.4 g) in acetonitrile (5 ml) was stirred at 60°C in the dark for 4 hours. The resulting pale pink solid was filtered off, and washed repeatedly with cold diethyl ether, to give 8-carbamoyl-3-(*p*-methoxybenzyl)-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one (0.23 g), m.p. 180-182°C (with effervescence).

EXAMPLE 21

By proceeding in a similar manner to the foregoing Examples, there was prepared 8-(N-allylcarbamoyl)-3-(2-chloroethyl)-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one [I.R. 1750 cm⁻¹; NMR (in DMSO-d₆): multiplets 3.96, 5.06 and 5.84 ppm: triplets 4.60 and 6.2 ppm: singlet 8.78 ppm], from 5-amino-4-allylcarbamoylimidazole via 5-diazo-4-allylcarbamoylimidazole.

The 5-amino-4-allylcarbamoylimidazole was prepared from 5-nitro-4-allylcarbamoylimidazole (m.p. 218-220°C) by reduction by means of titanous chloride.

REFERENCE EXAMPLE

(i) an intimate mixture of 5-nitroimidazole-4-carboxylic acid (2.0 g) and phosphorus pentachloride (2.67 g) was stirred and heated in an oil bath at 120°C for 1 hour. The resulting yellow slurry was evaporated at 60°C/0.1 mm Hg for 30 minutes, to give 1,6-dinitro-5*H*,10*H*-diimidazo[1,5-*a*:1',5'-*d*]pyrazine-5,10-dione (1.90 g) in the form of a yellow solid, m.p. 249-251°C (with decomposition). [ν_{\max} (KBr disc) 1750 cm⁻¹; m/e 278 (M⁺)].

Windaus, Ber., 1923, 56, 684 and Gireva, Chem. Abs. 59, 1622e, using the same method, describe their products as "5-nitroimidazole-4-carbonyl chloride".

(ii) Aqueous dimethylamine solution (25% w/v; 60 ml) was cooled to between 0° and 5°C and treated portionwise, with stirring, with 1,6-dinitro-5*H*,10*H*-diimidazo[1,5-*a*:1',5'-*d*]pyrazine-5,10-dione (6.0 g) in that temperature range. The resulting deep purple solution was stirred for 2 hours. The solution was evaporated at 50°C/10 mm Hg and then acidified by treatment with concentrated hydrochloric acid, to give an orange solution. This solution was extracted with ethyl acetate (7 × 200 ml), and the combined extracts were dried over magnesium sulphate, and evaporated, to give a yellow solid (6.6 g). This solid was triturated with toluene (50 ml) and then recrystallised from ethyl acetate, to give 5[4]-nitro-4[5]-dimethylcarbamoylimidazole (2.53 g), in the form of yellow crystals, m.p. 193-195°C. [Elemental analysis:—found: C,38.9; H,30.4%; C₆H₈N₄O₃ requires: C, 39.1; H,4.38; N,30.4%].

(iii) A solution of 5[4]-nitro-4[5]-dimethylcarbamoylimidazole (1.62 g) in dry dimethylformamide

shaken under hydrogen at atmospheric pressure and room temperature. After 3 hours, hydrogen absorption was complete (710 ml). The mixture was treated with charcoal and filtered through diatomaceous earth. The dark brown filtrate was evaporated at 50°C/0.1 mm Hg and the resulting residue was triturated with diethyl ether to give crude 5[4] - amino - 4[5] - dimethylcarbamoylimidazole (1.75 g), in the form of a dark brown crystalline solid, m.p.

179-181°C [ν_{\max} (KBR disc) 1595 cm^{-1} ; NMR in DMSO- d_6 : singlets at 3.2 and 7.0 δ], which was still contaminated with colloidal platinum and which was used in the next stage without further purification.

(iv) A stirred solution of sodium nitrite (0.79 g) in water (5.7 ml) was cooled to between 5° and 10°C and treated, dropwise, within this temperature range, with a solution of 5[4] - amino - 4[5] - dimethylcarbamoylimidazole (1.75 g) in aqueous acetic acid (1M; 17.6 ml) during 5 minutes. The resulting solution was extracted with ethyl acetate (4 x 40 ml), the combined extracts were dried over magnesium sulphate and evaporated at 30°C/10 mm Hg, to give 4[5] - diazo - 5[4] - dimethylcarbamoylimidazole (1.59 g), in the form of orange crystals, m.p. 101-103°C (with decomposition) [Elemental analysis:—found: C, 42.6; H, 4.17; N, 41.4%; $\text{C}_8\text{H}_8\text{N}_5\text{O}$ requires: C, 43.6; H, 4.27; N, 42.4%].

The present invention includes within its scope pharmaceutical compositions which comprise, as active ingredient, at least one tetrazine derivative of general formula I, together with a pharmaceutical carrier or coating. In clinical practice the compounds of general formula I will normally be administered orally, rectally, vaginally or parenterally, e.g. intravenously or intraperitoneally.

Methods of presentation of pharmaceutically active compounds are well known in the art and a suitable vehicle may be determined by the physician or pharmacist, depending upon such factors as the effect sought, the size, age, sex and condition of the patient and on the properties of the active compound. The compositions may also contain, as is usual in the art, such materials as solid or liquid diluents, wetting agents, preservatives, flavouring and colouring agents and the like.

Solid compositions for oral administration include compressed tablets, pills, dispersible powders, and granules. In such solid compositions one or more of the active compounds is, or are, admixed with at least one inert diluent such as calcium carbonate, potato starch, alginic acid, or lactose. The compositions may also comprise, as is normal practice, additional substances other than inert diluents, e.g. lubricating agents, such as magnesium stearate. Liquid compositions for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups and elixirs containing in part diluents commonly used in the art, such as water and liquid paraffin. Besides inert diluents such compositions may also comprise adjuvants, such as wetting and suspending agents, e.g. polyvinylpyrrolidone, and sweetening, flavouring, perfuming and preserving agents. The compositions according to the invention, for oral administration, also include capsules of absorbable material such as gelatin containing one

or more of the active substances with or without the addition of diluents or excipients.

Solid compositions for vaginal administration include pessaries formulated in manner known *per se* and containing one or more of the active compounds.

Solid compositions for rectal administration include suppositories formulated in manner known *per se* and containing one or more of the active compounds.

Preparation according to the invention for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, or emulsions. Examples of non-aqueous solvents or suspending media are polyethylene glycol, dimethyl sulphoxide, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. These compositions may also include adjuvants such as preserving, wetting, emulsifying and dispersing agents. They may be sterilised, for example, by filtration through a bacteria-retaining filter, by incorporation of sterilising agents in the compositions, or by irradiation. They may also be manufactured in the form of sterile solid compositions, which can be dissolved in sterile water or some other sterile injectable medium immediately before use.

The percentage of active ingredient in the compositions of the invention may be varied, it being necessary that it should constitute a proportion such that a suitable dosage for the therapeutic effect desired shall be obtained. Obviously several unit dosage forms may be administered at about the same time. In general, the preparations should normally contain at least 0.025% by weight of active substance when required for administration by injection; for oral administration the preparation will normally contain at least 0.1% by weight of active substance. The dose employed depends upon the desired therapeutic effect, the route of administration and the duration of the treatment.

The tetrazine derivatives of general formula I are useful in the treatment of malignant neoplasms, for example carcinomas, melanomas, sarcomas, lymphomas and leukaemias, at doses which are generally between 0.1 to 200, preferably between 1 and 20, mg/kg body weight per day.

The following Composition Examples illustrate pharmaceutical compositions according to the present invention.

COMPOSITION EXAMPLE 1

A solution suitable for parenteral administration was prepared from the following ingredients:—

8-Carbamoyl-3-(2-chloroethyl)-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one	1.0 g
Dimethyl sulphoxide	10 ml
Arachis oil	90 ml
by dissolving the 8-carbamoyl-3-(2-chloroethyl)-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one in the dimethyl sulphoxide and adding the arachis oil. The resulting solution was divided, under aseptic conditions, into ampoules at an amount of 10 ml per ampoule. The ampoules were sealed, to give 10 ampoules each containing 100 mg of 8-carbamoyl-3-(2-chloroethyl)-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one.	

Similar ampoules containing solutions suitable for parenteral administration may be prepared by proceeding in a similar manner but replacing the 8-carbamoyl-3-(2-chloroethyl)-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one by another compound of general formula I.

COMPOSITION EXAMPLE 2


Capsules suitable for oral administration were prepared by placing 8-carbamoyl-3-(2-chloroethyl)-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one into gelatin shells of number 2 size at a rate of 10 mg per capsule.

Similar capsules may be prepared by using another compound of general formula I or any other conveniently sized capsule shells.

CLAIMS

1. [3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one derivatives of the general formula:-

20



25 wherein R¹ represents a hydrogen atom, or a straight- or branched-chain alkyl, alkenyl or alkynyl group containing up to 6 carbon atoms, each such group being unsubstituted or substituted by from one to three substituents selected from halogen atoms, straight- or branched-chain alkoxy, alkylthio, alkylsulphinyl and alkylsulphonyl groups containing up to 4 carbon atoms, and optionally substituted phenyl groups, or R¹ represents a cycloalkyl group containing from 3 to 8 carbon atoms, and R² represents a carbamoyl group which may carry on the nitrogen atom one or two groups selected from straight- and branched-chain alkyl and alkenyl groups, each containing up to 4 carbon atoms, and cycloalkyl groups containing from 3 to 8 carbon atoms and - when R¹ represents hydrogen - alkali metal salts thereof.

2. Tetrazine derivatives according to claim 1 wherein R¹ represents an alkyl, alkenyl or alkynyl group substituted by one, two or three optionally substituted phenyl groups and the optional substituents on the phenyl radical(s) are selected from alkoxy and alkyl groups containing up to 4 carbon atoms, and the nitro group.

3. Tetrazine derivatives according to claim 1 wherein R¹ represents a straight- or branched-chain alkyl group containing from 1 to 6 carbon atoms optionally substituted by one or two halogen atoms or by an alkoxy group containing 1 to 4 carbon atoms or by a phenyl group optionally substituted by one or two alkoxy groups containing from 1 to 4 carbon atoms, or R¹ represents an alkenyl group containing 2 to 6 carbon atoms or a cyclohexyl group.

4. Tetrazine derivatives according to claim 3 wherein the halogen atom(s) is (or are) chlorine, fluorine and/or bromine, the alkoxy group(s) is (or are) methoxy, and the alkenyl group is allyl.

5. Tetrazine derivatives according to claim 1 wherein R¹ represents a straight- or branched-chain alkyl group containing from 1 to 6 carbon atoms, unsubstituted or substituted by a halogen atom

6. Tetrazine derivatives according to claim 5 wherein R¹ represents an alkyl group containing 1 to 3 carbon atoms unsubstituted or substituted by a halogen atom.

7. Tetrazine derivatives according to claim 1 or 5 wherein R¹ represents a methyl or 2-haloalkyl group.

8. Tetrazine derivatives according to claim 5, 6 or 7 in which the halogen atom on the alkyl group is chlorine or fluorine.

9. Tetrazine derivatives according to claim 1 wherein R¹ represents a 2-fluoroethyl or 2-chloroethyl group.

10. Tetrazine derivatives according to claim 1 wherein R¹ represents a benzyl or *p*-methoxybenzyl group.

11. Tetrazine derivatives according to any one of claims 1 to 10 wherein R² represents a carbamoyl group, a monoalkylcarbamoyl group containing up to 4 carbon atoms in the alkyl radical, or a monoalkenylcarbamoyl group containing up to 4 carbon atoms in the alkenyl radical.

12. Tetrazine derivatives according to claim 1 wherein R¹ represents a straight- or branched-chain alkyl, alkenyl or alkynyl group containing from 1 to 6 carbon atoms, each such group being unsubstituted or substituted by from one to three halogen atoms, and R² represents the carbamoyl group.

13. 8-Carbamoyl-3-methyl-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one.

14. 8-Carbamoyl-3-n-propyl-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one.

15. 8-Carbamoyl-3-(2-chloroethyl)-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one.

16. 3-(2-Chloroethyl)-8-methylcarbamoyl-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one.

17. 8-Carbamoyl-3-(3-chloropropyl)-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one.

18. 8-Carbamoyl-3-(2,3-dichloropropyl)-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one.

19. 3-Allyl-8-carbamoyl-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one.

20. 3-(2-Chloroethyl)-8-dimethylcarbamoyl-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one.

21. 3-(2-Bromoethyl)-8-carbamoyl-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one.

22. 3-Benzyl-8-carbamoyl-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one.

23. 8-Carbamoyl-3-(2-methoxyethyl)-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one.


24. 8-Carbamoyl-3-cyclohexyl-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one.

25. 8-Carbamoyl-3-(*p*-methoxybenzyl)-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one.

26. 8-(N-Allylcarbamoyl)-3-(2-chloroethyl)-[3H]-imidazo[5,1-d]-1,2,3,5-tetrazin-4-one.

27. A process for the preparation of a tetrazine derivative of the general formula depicted in claim 1, wherein R¹ and R² are as defined in claim 1 but excluding hydrogen from the definition of R¹, which comprises reacting a compound of the general formula:

130




II

(wherein R² is as defined in claim 1) with an isocyanate of the general formula:—




- 5 wherein R³ represents an alkyl, alkenyl or alkynyl group containing up to 6 carbon atoms, each such group being unsubstituted or substituted by one to three substituents selected from halogen atoms, alkoxy, alkylthio, alkylsulphinyl and alkylsulphonyl groups containing up to 4 carbon atoms, and optionally substituted phenyl groups, or R³ represents a cycloalkyl group containing 3 to 8 carbon atoms.
- 10 28. A process for the preparation of a tetrazine derivative of the general formula depicted in claim 1, wherein R¹ and R² are as defined in claim 1 but excluding hydrogen from the definition of R¹, which comprises reacting a compound of the general formula:
- 20



- 25 (wherein R² is as defined in claim 1) or an alkali metal salt thereof with a compound of the general formula:—
- 30




- wherein R³ is as defined in claim 27, and X represents the acid residue of a reactive ester.
- 35 29. A process according to claim 28 wherein X represents a halogen atom or a methoxysulphonyloxy, methanesulphonyloxy or toluene-*p*-sulphonyloxy group.
30. A process for the preparation of a tetrazine derivative of the general formula depicted in claim 1, wherein R¹ represents a hydrogen atom and R² is as defined in claim 1, or an alkali metal salt thereof which comprises reacting a compound of the general formula:
- 40



- 45 with a compound of the general formula:—
- 50



- wherein R⁴ represents an alkali metal atom or a protecting group such as a benzyl or *p*-methoxybenzyl group, followed, when R⁴ represents a protecting group, by the replacement of the protecting group by a hydrogen atom in the compound thus obtained of the general formula:—
- 55



60

wherein R² is as defined in claim 1 and R⁵ represents a protecting group such as a benzyl or *p*-methoxybenzyl group, by methods known *per se*.

31. A process for the preparation of a tetrazine derivative of the general formula depicted in claim 1, wherein R¹ and R² are as defined in claim 1, or — when R¹ represents hydrogen — an alkali metal salt thereof substantially as hereinbefore described with especial reference to any of Examples 1 to 21.
- 70 32. Pharmaceutical compositions which comprise, as active ingredient, at least one tetrazine derivative as claimed in any one of claims 1 to 26 in association with a pharmaceutical carrier or coating.
- 75 33. Pharmaceutical compositions according to claim 32 substantially as hereinbefore described with especial reference to Composition Example 1 to 2.
- 80 34. Tetrazine derivatives of the general formula depicted in claim 1, wherein R¹ and R² are as defined in claim 1, for use in the treatment of malignant neoplasms such as carcinomas, melanomas, sarcomas, lymphomas and leukaemias.
- 85 35. Tetrazine derivatives of the general formula depicted in claim 1, wherein R¹ and R² are as defined in claim 1, for use in the treatment of organ grafts and skin grafts and in the treatment of immunological diseases.
- 90

Printed for Her Majesty's Stationery Office by The Tweeddale Press Ltd., Berwick-upon-Tweed, 1983.

Published at the Patent Office, 25 Southampton Buildings, London, WC2A 1AY, from which copies may be obtained.